

AD-A069 290

STANFORD UNIV CALIF DEPT OF OPERATIONS RESEARCH
INSCRIBING AND CIRCUMSCRIBING CONVEX POLYHEDRA.(U)
MAR 79 B C EAVES, R M FREUND

F/6 12/1

UNCLASSIFIED

ARO-15254.6-M

DAAG29-78-G-0026


NL

1 OF 1
AD
A069290



END
DATE
FILMED

7-79
DDC



NATIONAL BUREAU OF STANDARDS
MICROCOPY RESOLUTION TEST CHART

DDC FILE COPY

ADA069290

(18) AR8

(19) 152 54.6-M

(6) INSCRIBING AND CIRCUMSCRIBING CONVEX POLYHEDRA .

by

(10) B. Curtis / Eaves ~~and~~ R. M. / Freund

(9) TECHNICAL REPORT ,

(11) March 1979

(12) 14 p.

Department of Operations Research
Stanford University
Stanford, California

DDC
RECEIVED
JUN 4 1979
A

DISTRIBUTION STATEMENT A
Approved for public release;
Distribution Unlimited

(15)
This research was sponsored in part by the Army Research Office -
Durham, Contract No. DAAG 29-78-G-0026, and the National Science
Foundation, MCS 77-05623

✓NSF

402 766
79 05 29 024
Liu


TABLE OF CONTENTS

Chapter		Page No.
1.	Introduction	1
2.	Preliminaries	3
3.	Results	6
4.	Related Problems	10

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/ _____	
Availability Codes	
Dist.	Avalland/or special
A	

1. Introduction

Let \mathcal{X} and \mathcal{Y} be polyhedra; that is, closed polyhedral convex sets, bounded or not, in \mathbb{R}^n . Our interest is in computing the smallest nonnegative scale $s\mathcal{X}$ of \mathcal{X} for which some translate $s\mathcal{X} + t$ contains \mathcal{Y} , or equivalently, of computing the largest nonnegative scale $s\mathcal{Y}$ of \mathcal{Y} for which some translate $s\mathcal{Y} + t$ is contained in \mathcal{X} .



For certain descriptions of \mathcal{X} and \mathcal{Y} we observe that this problem; namely, the circumscription program

$$P1 \quad \begin{cases} z_1 = \inf_{s,t} s \\ \text{subject to: } \mathcal{Y} \subseteq s\mathcal{X} + t \quad s \geq 0 \end{cases}$$

or equivalently, the inscription program

$$P2 \quad \begin{cases} z_2 = \sup_{s,t} s \\ \text{subject to: } s\mathcal{Y} + t \subseteq \mathcal{X} \quad s \geq 0 \end{cases}$$

is a linear program.

That an n -sphere of maximum radius in a polyhedron can be found by linear programming has been in the folklore for over a decade and has been used in a variety of applications. Perhaps our observations will be of use as well.

2. Preliminaries

To avoid trivialities we assume throughout that \mathcal{X} and \mathcal{Y} contain at least one and two points, respectively.

\mathcal{X} and \mathcal{Y} can be described as the intersection of finitely many hyperplanes

$$\mathcal{X} = \{x \in \mathbb{R}^n : Ax \leq a\} \quad \mathcal{Y} = \{x \in \mathbb{R}^n : Bx \leq b\}$$

in which case we say that \mathcal{X} and \mathcal{Y} have representation H .

Alternatively, \mathcal{X} and \mathcal{Y} can be described as a weighting of points and rays

$$\mathcal{X} = \{x \in \mathbb{R}^n : x = X\lambda + U\omega, e\lambda = 1, \lambda \geq 0, \omega \geq 0\}$$

$$\mathcal{Y} = \{x \in \mathbb{R}^n : x = Y\mu + V\pi, e\mu = 1, \mu \geq 0, \pi \geq 0\}$$

where $e = (1, 1, \dots, 1)$ in which case we say \mathcal{X} and \mathcal{Y} have representation W .

Given a representation one can systematically convert it to the other; however, we shall suppose, and typically rightly so, that the computational burden of this conversion is prohibitive.

Using the representation W we see that any polyhedron \mathcal{X} can be expressed as $K + C$ where K is a compact polyhedron and C is the polyhedron of rays of \mathcal{X} . For any positive scale s we

have $s\mathcal{K} = sK + sC = sK + C$, and, of course, for a zero scale s we have $s\mathcal{K} = \{0\}$. Thus the set $s\mathcal{K}$ is continuous in s for positive s , and is continuous in nonnegative s if and only if C contains only the origin.

Given \mathcal{K} the cone of rays C is unique and we write $C = \text{cone } \mathcal{K}$. By $\text{tng } \mathcal{K}$ we mean the smallest subspace of R^n that contains a translate of \mathcal{K} .

The next lemmas describe the sense in which $P1$ and $P2$ are equivalent.

Lemma 1 (Principal Equivalence): If (s_1, t_1) is feasible or optimal for $P1$ and s_1 is positive, then $(s_2, t_2) = (1/s_1, -t_1/s_1)$ is feasible or optimal, respectively, for $P2$, and vice versa. \square

Lemma 2 (Feasibility): $P1$ is feasible if and only if $\text{cone } \mathcal{Y} \subseteq \text{cone } \mathcal{K}$ and $\text{tng } \mathcal{Y} \subseteq \text{tng } \mathcal{K}$. $P2$ is always feasible. \square

Lemma 3 (Attainment): The following are equivalent.

- | | |
|--------------------------|------------------------------------|
| i) $0 < z_1 < +\infty$ | ii) $P1$ has an optimum |
| iii) $0 < z_2 < +\infty$ | iv) $P2$ has an optimum. \square |

Lemma 4 (Non-attainment): The following are equivalent:

- i) $z_1 = 0$
- ii) $z_2 = +\infty$
- iii) $\mathcal{V} + t \subseteq \text{cone } \mathcal{X}$ for some t □

The possible discontinuity of $s\mathcal{X} + t$ at $s = 0$
accounts for the incomplete equivalence between P1 and P2 .

3. Results

In this section we formulate the circumscription and inscription problems P1 and P2 for three cases of representation; namely, HH, WW, and HW, where, for example, HW refers to \mathcal{X} and \mathcal{Y} having representations H and W, respectively.

In each case a more general problem is treated first. We generalize P1 and P2 to

$$P3 \begin{cases} \text{infimum: } c\theta \\ \theta \\ \text{subject to: } \mathcal{Y} \subseteq \mathcal{X}(\theta) \end{cases} \quad \theta \in \Theta$$

and

$$P4 \begin{cases} \text{supremum: } c\theta \\ \theta \\ \text{subject to: } \mathcal{Y}(\theta) \subseteq \mathcal{X} \end{cases} \quad \theta \in \Theta$$

where Θ is a polyhedron in R^k and $c\theta$ is a linear function of θ in Θ that measures some feature of the polyhedrons. We regain P1 and P2 from P3 and P4 by setting $\theta = (s,t)$, etc.

Case HH

Let $\mathcal{X}(\theta)$ be the set $\{x : A(\theta)x \leq a(\theta)\}$ where (A,a) is an affine function of θ in Θ and let $\mathcal{Y} = \{x : Bx \leq b\}$. The program P3 is seen to be, using an alternative theorem, the linear program

$$P5 \quad \begin{cases} \text{minimize: } c\theta \\ \theta, \Lambda \\ \text{subject to: } \Lambda B = A(\theta) & \Lambda b \leq a(\theta) \\ \Lambda \geq 0 & \theta \in \Theta \end{cases}$$

Observing that $s\mathcal{X} + t = \{x : Ax \leq sa + At\}$ for $s > 0$ and specializing P5 to solve P1 we obtain the linear program

$$P6 \quad \begin{cases} z_1 = \text{minimize: } s \\ s, t, \Lambda \\ \text{subject to: } \Lambda B = A & \Lambda b \leq as + At \\ \Lambda \geq 0 & s \geq 0 \end{cases}$$

Case WW

Let $\mathcal{Y}(\theta)$ be the set $\{Y(\theta)\mu + V(\theta)\pi : e\mu = 1, \mu \geq 0, \pi \geq 0\}$ where (Y, V) is an affine function of θ in Θ and let \mathcal{X} be the set $\{X\lambda + U\omega : e\lambda = 1, \lambda \geq 0, \omega \geq 0\}$. The program P4 is the linear program

$$P7 \quad \begin{cases} \text{maximize: } c\theta \\ \theta, \Lambda, \Omega, \Pi \\ \text{subject to: } Y(\theta) = X\Lambda + U\Omega & e\Lambda = e \\ V(\theta) = U\Pi & \theta \in \Theta \\ \Lambda \geq 0 & \Omega \geq 0 & \Pi \geq 0 \end{cases}$$

Specializing P7 to solve P2 we obtain the linear program

$$P8 \left\{ \begin{array}{ll} z_2 = \text{maximize:} & s \\ & s, t, \Lambda, \Omega, \Pi \\ \text{subject to:} & t \circ e + sY = X\Lambda + U\Omega \quad e\Lambda = e \\ & sV = U\Pi \quad s \geq 0 \\ & \Lambda \geq 0 \quad \Omega \geq 0 \quad \Pi \geq 0 \end{array} \right.$$

where \circ denotes outer product. In solving P8 one first verifies that $V = U\Pi$ with $\Pi \geq 0$ has a solution, and then drops the constraints $sV = U\Pi$ and $\Pi \geq 0$.

Case HW₁

We treat the case HW twice as HW₁ and HW₂ where we approach the circumscription/inscription problems through P3 and P4, respectively. Let $\mathcal{X}(\theta) = \{x : A(\theta)x \leq a(\theta)\}$ where (A, a) is an affine function of θ in Θ and let $\mathcal{Y} = \{Y\mu + V\pi : e\mu = 1, \mu \geq 0, \pi \geq 0\}$. The program P3 is the linear program

$$P9 \left\{ \begin{array}{ll} \text{minimize:} & c\theta \\ & \theta \\ \text{subject to:} & A(\theta)Y \leq a(\theta) \circ e \\ & A(\theta)V \leq 0 \quad \theta \in \Theta \end{array} \right.$$

Specializing P9 to solve P1 we obtain the linear program

$$P10 \quad \left\{ \begin{array}{ll} z_1 = \text{minimize:} & s \\ & s, t \\ \text{subject to:} & AY \leq saoe + A(Toe) \\ & AV \leq 0 \quad s \geq 0 \end{array} \right.$$

In solving P10 one verifies $AV \leq 0$ and then drops the constraints $AV \leq 0$.

Case HW₂

Let $\mathcal{Y}(\theta)$ be the set $\{Y(\theta)\mu + V(\theta)\pi : e\mu = 1, \mu \geq 0, \pi \geq 0\}$ where (Y, V) is an affine function of θ in Θ and \mathcal{X} is the set $\{x : Ax \leq a\}$. Then the program P4 is the linear program

$$P11 \quad \left\{ \begin{array}{ll} \text{maximize:} & c\theta \\ & \theta \\ \text{subject to:} & AY(\theta) \leq aoe \\ & AV(\theta) \leq 0 \quad \theta \in \Theta \end{array} \right.$$

Specializing P11 to solve P2 we obtain the linear program

$$P12 \quad \left\{ \begin{array}{ll} z_2 = \text{maximize:} & s \\ & s, t \\ \text{subject to:} & A(sY + Toe) \leq a \\ & sAV \leq 0 \quad s \geq 0 \end{array} \right.$$

In solving P12 one verifies $AV \leq 0$ and then drops the constraints $sAV \leq 0$. Observe that P11 remains a linear program if a also is an affine function of θ in Θ .

4. Related Problems

The forgoing raises the question as to whether the following problems can be solved as linear programs.

- a) Case WH
- b) Finding the largest \mathcal{Y} in \mathcal{X} or smallest \mathcal{X} containing \mathcal{Y} where rotations as well as scales and translations are permitted.
- c) Finding the largest n -sphere \mathcal{Y} in \mathcal{X} where \mathcal{X} has representation W .
- d) Finding the smallest n -sphere \mathcal{X} containing \mathcal{Y} where \mathcal{Y} has representation H or W .

We suspect, but have no comprehensive proofs, that none of these problems can be formulated as a linear program. Observe in a) that for fixed (X, U, B, b) Case WH can be formulated as a linear program by converting to one of the other cases; in b) the set of optimal \mathcal{Y} may not be a convex set; and in d) the n -sphere may have an irrational radius given rational data.

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) INSCRIBING AND CIRCUMSCRIBING CONVEX POLYHEDRA		5. TYPE OF REPORT & PERIOD COVERED Technical Report
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) B. Curtis Eaves and R. M. Freund		8. CONTRACT OR GRANT NUMBER(s) DAAG-29-78-G-0026
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Operations Research Stanford University Stanford, California 94305		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS Mathematics Division U.S. Army Research Office Box CM, Duke Station, Durham, N.C. 27706		12. REPORT DATE March 1979
		13. NUMBER OF PAGES 13
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE NA
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) NA		
18. SUPPLEMENTARY NOTES The findings of this report are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Inscribing Circumscribing Polyhedra Linear Programming		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Let \mathcal{X} and \mathcal{Y} be closed polyhedral convex sets, bounded or not, in \mathbb{R}^n . For certain representations of \mathcal{X} and \mathcal{Y} it is shown that the task of finding the smallest scale of \mathcal{X} for which some translate contains \mathcal{Y} can be resolved with linear programming.		

DD FORM 1473
1 JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-014-6601

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)